arXiv:1408.4980v1 [math.NT] 21 Aug 2014

FAMILY COMPLEXITY AND CROSS-CORRELATION MEASURE FOR FAMILIES OF BINARY SEQUENCES

ARNE WINTERHOF AND OĞUZ YAYLA

Abstract. We study the relationship between two measures of pseudorandomness for families of binary sequences: family complexity and cross-correlation measure introduced by Ahlswede et al. in 2003 and recently by Gyarmati et al., respectively. More precisely, we estimate the family complexity of a family $(e_{i,1},\ldots,e_{i,N}) \in \{-1,+1\}^N$, $i=1,\ldots,F$, of binary sequences of length N in terms of the cross-correlation measure of its dual family $(e_{1,n},\ldots,e_{F,n}) \in \{-1,+1\}^F$, $n=1,\ldots,N$.

We apply this result to the family of sequences of Legendre symbols with irreducible quadratic polynomials modulo p with middle coefficient 0, that is, $e_{i,n} = \left(\frac{n^2-bi^2}{p}\right)_{n=1}^{(p-1)/2}$ for $i=1,\ldots,(p-1)/2$, where b is a quadratic nonresidue modulo p, showing that this family as well as its dual family have both a large family complexity and a small cross-correlation measure up to a rather large order.

1. Introduction

Pseudorandom binary sequences are used in many areas such as telecommunication, cryptography, simulation, spectroscopy, see for example [2, 6, 11]. The quality of a pseudorandom sequence has to be screened by statistical test packages (for example L'Ecuyer's TESTU01, Marsaglia's Diehard or the NIST battery) as well as by proving theoretical results on certain measures of pseudorandomness such as the correlation measure of order ℓ introduced by Mauduit and Sárközy [9]. Here we focus on theoretical results.

In many applications such as cryptography one needs a large family of good pseudorandom sequences and has to prove bounds on several figures of merit [6, 7]. In this paper we study the relationship of two such quality measures, the family complexity, short f-complexity, and the cross-correlation measure of order ℓ of families of binary sequences.

Ahlswede et al. [1] introduced the f-complexity as follows.

Definition 1.1. The *f-complexity* $C(\mathcal{F})$ of a family $\mathcal{F}$ of binary sequences $E_N \in \{-1,+1\}^N$ of length N is the greatest integer $j \geq 0$ such that for any $1 \leq i_1 < i_2 < \cdots < i_j \leq N$ and any $\epsilon_1,\epsilon_2,\ldots,\epsilon_j \in \{-1,+1\}$ there is a sequence $E_N = \{e_1,e_2,\ldots,e_N\} \in \mathcal{F}$ with

$$e_{i_1} = \epsilon_1, e_{i_2} = \epsilon_2, \ldots, e_{i_j} = \epsilon_j.$$

2010 *Mathematics Subject Classification.* 11K45, 11T24.

Key words and phrases. pseudorandomness, binary sequences, family complexity, cross-correlation measure, Legendre sequence, polynomials over finite fields.

We have the trivial upper bound

$$2^{C(\mathcal{F})} \le |\mathcal{F}|, \tag{1}$$

where $|\mathcal{F}|$ denotes the size of the family $\mathcal{F}$.

Gyarmati et al. [7] introduced the cross-correlation measure of order ℓ.

Definition 1.2. The *cross-correlation measure of order* ℓ of a family $\mathcal{F}$ of binary sequences $E_{i,N} = (e_{i,1}, e_{i,2}, \ldots, e_{i,N}) \in \{-1+1\}^N$, $i = 1, 2, \ldots, F$, is defined as

$$\Phi_\ell(\mathcal{F}) = \max_{M,D,I} \left| \sum_{n=1}^{M} e_{i_1,n+d_1} \cdots e_{i_\ell,n+d_\ell} \right|,$$

where D denotes an ℓ tuple $(d_1, d_2, \ldots, d_\ell)$ of integers such that $0 \le d_1 \le d_2 \le \cdots \le d_\ell < M + d_\ell \le N$ and $d_i \ne d_j$ if $E_{i,N} = E_{j,N}$ for $i \ne j$, and I denotes an ℓ tuple $(i_1, i_2, \ldots, i_\ell) \in \{1, 2, \ldots, F\}^\ell$.

In Section 2 we estimate the f-complexity of a family of binary sequences

$$E_{i,N} = (e_{i,1}, e_{i,2}, \ldots, e_{i,N}) \in \{-1+1\}^N, \quad i = 1, \ldots, F,$$

in terms of the cross-correlation measure of the *dual family* $\overline{\mathcal{F}}$ of binary sequences

$$E_{i,F} = (e_{1,i}, e_{2,i}, \ldots, e_{F,i}) \in \{-1+1\}^F, \quad i = 1, \ldots, N.$$

In Section 3 we apply this result to prove a bound on the f-complexity of the family of sequences of Legendre-symbols of monic quadratic irreducible polynomials with vanishing middle coefficient

$$\mathcal{F} = \left\{ \left(\frac{n^2 - bi^2}{p} \right)_{n=1}^{(p-1)/2} : i = 1, \ldots, (p-1)/2 \right\},$$

where b is a quadratic non-residue modulo p, showing that this family as well as its dual family have both a large family complexity and a small cross-correlation measure up to a rather large order.

We note that there are several related constructions of families of binary sequences defined with the Legendre symbol and polynomials, see [5, 7] and references therein. For instance, the family given in [5] has a large family complexity but a large cross-correlation measure of order 2. Moreover, the families given in [7] have a small cross-correlation measure but it is not easy to measure their family complexity, for further details see the remarks in Section 3. One of the aims of this study is to construct a family of binary sequences having both a large family complexity and a small cross-correlation measure.

Throughout the paper, the notation $U \ll V$ is equivalent to the statement that $|U| \le cV$ holds with some positive constant c. Moreover, the notation $f(n) = o(1)$ is equivalent to

$$\lim_{n\to\infty} f(n) = 0.$$

2. A relation between family complexity and cross-correlation measure

In this section we prove the following relationship between the f-complexity of a family of binary sequences and the cross-correlation measure of its dual family.

Theorem 2.1. *Let $\mathcal{F}$ be a family of binary sequences $(e_{k,1},\ldots,e_{k,N}) \in \{-1,+1\}^N$ for $k = 1,2,\ldots,F$ and $\overline{\mathcal{F}}$ its dual family of binary sequences $(e_{1,n},e_{2,n},\ldots,e_{F,n}) \in \{-1,+1\}^F$ for $n = 1,2,\ldots,N$. Then we have*

$$C(\mathcal{F}) \geq \left\lceil \log_2 F - \log_2 \max_{1\leq i\leq \log_2 F} \Phi_i(\overline{\mathcal{F}}) \right\rceil - 1,$$

where $\log_2$ denotes the binary logarithm.

Proof. We are looking for the largest j such that any specification

$$e_{k,n_1} = \epsilon_1, e_{k,n_2} = \epsilon_2, \ldots, e_{k,n_j} = \epsilon_j \tag{2}$$

occurs in the family $\mathcal{F}$ for some $k \in \{1,2,\ldots,F\}$. We will prove a sufficient condition for the existence of a sequence in $\mathcal{F}$ satisfying (2) by a counting argument. We know that

$$\frac{1}{2}(1+\epsilon_i e_{k,n_i}) = \begin{cases} 1 & \text{if } e_{k,n_i} = \epsilon_i, \\ 0 & \text{if } e_{k,n_i} = -\epsilon_i. \end{cases}$$

Then the number A of sequences in $\mathcal{F}$ satisfying (2) equals

$$A = \sum_{k=1}^{F} \frac{1}{2^j} \prod_{i=1}^{j} (1+\epsilon_i e_{k,n_i}).$$

Extending the product and after easy calculations we have

$$\begin{aligned} A &= \frac{1}{2^j} \sum_{k=1}^{F} \left[1 + \sum_{\ell=1}^{j} \sum_{1\leq i_1<\cdots<i_\ell\leq j} \epsilon_{i_1}\cdots\epsilon_{i_\ell} e_{k,n_{i_1}}\cdots e_{k,n_{i_\ell}} \right] \\ &= \frac{1}{2^j} \left[F + \sum_{\ell=1}^{j} \sum_{1\leq i_1<\cdots<i_\ell\leq j} \epsilon_{i_1}\cdots\epsilon_{i_\ell} \sum_{k=1}^{F} e_{k,n_{i_1}}\cdots e_{k,n_{i_\ell}} \right] \\ &\geq \frac{1}{2^j} \left[F - \sum_{\ell=1}^{j} \sum_{1\leq i_1<\cdots<i_\ell\leq j} \left| \sum_{k=1}^{F} e_{k,n_{i_1}}\cdots e_{k,n_{i_\ell}} \right| \right] \\ &\geq \frac{1}{2^j} \left[F - \sum_{\ell=1}^{j} \binom{j}{\ell} \Phi_\ell(\overline{\mathcal{F}}) \right]. \end{aligned}$$

Thus if

$$F > \sum_{\ell=1}^{j} \binom{j}{\ell} \Phi_\ell(\overline{\mathcal{F}}), \tag{3}$$

then there exists at least one sequence in $\mathcal{F}$ satisfying (2).

By (1) we may assume $j \leq \log_2 F$, and so we get

$$\sum_{\ell=1}^{j} \binom{j}{\ell} \Phi_\ell(\overline{\mathcal{F}}) \quad \leq \quad 2^j \max_{1\leq i\leq \log_2 F} \Phi_i(\overline{\mathcal{F}}).$$

Therefore for all integers $j \geq 0$ satisfying

$$j < \log_2 F - \log_2 \max_{1 \leq \ell \leq \log_2 F} \Phi_\ell(\overline{\mathcal{F}})$$

(3) is satisfied and we have $A > 0$ which completes the proof. □

3. A family with low cross-correlation measure and high family complexity

In this section we demonstrate how to apply Theorem 2.1 and prove that the following family of sequences and its dual family have both high family complexity and small cross-correlation measure of order ℓ up to a large order ℓ.

Let $p > 2$ be a prime and b a quadratic nonresidue modulo p. The family $\mathcal{F}$ and its dual family $\overline{\mathcal{F}}$ are defined as follows:

$$\mathcal{F} = \left\{ \left(\frac{n^2 - bi^2}{p} \right)_{i=1}^{(p-1)/2} : n = 1, \ldots, (p-1)/2 \right\},$$

$$\overline{\mathcal{F}} = \left\{ \left(\frac{n^2 - bi^2}{p} \right)_{n=1}^{(p-1)/2} : i = 1, \ldots, (p-1)/2 \right\}.$$

Since $\left(\frac{n^2-bi^2}{p}\right) = \left(\frac{-b}{p}\right)\left(\frac{i^2-bn^2}{p}\right) = (-1)^{(p+1)/2}\left(\frac{i^2-bn^2}{p}\right)$ both families have the same family complexity and cross-correlation measure of order ℓ. We will show

$$Q_\ell(\mathcal{F}) \ll \ell p^{1/2} \log p \quad \text{and} \quad Q_\ell(\overline{\mathcal{F}}) \ll \ell p^{1/2} \log p \tag{4}$$

for each integer $\ell = 1, 2, \ldots$ Then Theorem 2.1 immediately implies

$$C(\mathcal{F}) \geq \left(\frac{1}{2} - o(1) \right) \frac{\log p}{\log 2} \quad \text{and} \quad C(\overline{\mathcal{F}}) \geq \left(\frac{1}{2} - o(1) \right) \frac{\log p}{\log 2}.$$

Here we used

$$|\mathcal{F}| = |\overline{\mathcal{F}}| = (p-1)/2, \quad p \geq 11$$

which can be verified in the following way. For some $1 \leq n_1 < n_2 \leq (p-1)/2$ assume that $\left(\frac{n_1^2-bi^2}{p}\right) = \left(\frac{n_2^2-bi^2}{p}\right)$ for $i = 1, \ldots, (p-1)/2$. Since $i^2 \equiv (p-i)^2 \bmod p$ and $\left(\frac{n_1^2}{p}\right) = \left(\frac{n_2^2}{p}\right) = 1$ these Legendre symbols are the same for all $i = 0, \ldots, p-1$ and thus

$$p = \sum_{i=0}^{p-1} \left(\frac{n_1^2 - bi^2}{p} \right) \left(\frac{n_2^2 - bi^2}{p} \right) \leq 3p^{1/2}$$

by Weil's bound and we get a contradiction to $p \geq 11$.

Now we prove (4). According to Definition 1.2 and since the Legendre symbol is multiplicative we need to estimate sums of the form

$$\left| \sum_{n=1}^{M} \left(\frac{h(n)}{p} \right) \right|$$

where

$$h(X) = ((X + d_1)^2 - bi_1^2)((X + d_2)^2 - bi_2^2) \cdots ((X + d_\ell)^2 - bi_\ell^2).$$

The result follows from the Weil bound after reducing these incomplete character sums to complete ones using the standard method provided that $h(X)$ is not a square, see [8, 10, 12]. Since each factor of $h(X)$ of the form $(X + d)^2 - bi^2$ is

irreducible over $\mathbb{F}_p$, it is enough to show that one factor is distinct from all the others. We may assume that the first factor is of the form $X^2 - bi^2$. Comparing coefficients we see that

$$X^2 - bi^2 = (X + d)^2 - bj^2$$

is only possible if $d = 0$ and $i \equiv \pm j \bmod p$. Since $1 \le i, j \le (p-1)/2$ we get $i = j$ and $h(X)$ is not a square since either $d_j \ne d_k$ for $i \ne k$ or $d_j = d_k$ and $i_j \ne i_k$ with $1 \le i_j, i_k \le (p-1)/2$.

Remarks.

1. Gyarmati [5] studied the family $\mathcal{F}$ of binary sequences $E_f = (e_{f,1}, \dots, e_{f,p}) \in \{-1, +1\}^p$ defined by

$$e_{f,n} = \begin{cases} \left(\frac{f(n)}{p}\right) & \text{if } \gcd(f(n), p) = 1, \\ 1 & \text{if } p \mid f(n), \end{cases} \tag{5}$$

   where f runs through all non-constant square-free polynomials over $\mathbb{F}_p$ of degree at most k. She proved

$$C(F) \ge \frac{K}{2 \log 2} \log p - O(k \log (k \log p)).$$

   However, this family has obviously a very large cross-correlation measure of order 2. (Compare E_f and E_g with $g(X) = f(X + d)$ for some $d \in \mathbb{F}_p^*$.)
2. Gyarmati, Mauduit, and Sárközy [3] studied the family $\mathcal{F}$ of sequences $E_f = (e_{f,1}, \dots, e_{f,p}) \in \{-1, +1\}^p$ defined by (5) where f runs through all monic irreducible polynomials of degree d with second leading coefficient 0 and showed in Theorem 8.14 that

$$\Phi_l(\mathcal{F}) \ll \ell d p^{1/2} \log p.$$

   We note that for $d = 2$ the polynomials f are of the form $X^2 - a$ where a is a quadratic non-residue modulo p and we have $e_{f,1} = e_{f,p-1}$ and thus $C(\mathcal{F}) \le 1$. For $d \ge 3$ it seems to be a challenging problem to estimate the family complexity of this family.
3. Very recently, Gyarmati [4] proved also a lower bound on the family complexity of the family of sequences E_f defined by (5) where f runs through *all* monic irreducible polynomials of degree d (with arbitrary second leading coefficient). However, since $f(X)$ is irreducible whenever $f(X + c)$ for all $c \in \mathbb{F}_p$ the cross-correlation measure of order 2 of this family is again very large.

Acknowledgment

The first author is supported by the Austrian Science Fund (FWF): Project F5511-N26 which is part of the Special Research Program "Quasi-Monte Carlo Methods: Theory and Applications". The second author is supported by TÜBİTAK under Grant No. 2219.

References

1. Rudolf Ahlswede, Levon H. Khachatrian, Christian Mauduit, and András Sárközy, *A complexity measure for families of binary sequences*, Period. Math. Hungar. **46** (2003), no. 2, 107–118. MR 2004667 (2004j:11085)
2. Solomon W. Golomb and Guang Gong, *Signal design for good correlation*, Cambridge University Press, Cambridge, 2005. MR 2156522 (2006d:94021)
3. Louis Goubin, Christian Mauduit, and András Sárközy, *Construction of large families of pseudorandom binary sequences*, J. Number Theory **106** (2004), no. 1, 56–69. MR 2049592 (2004m:11121)
4. Katalin Gyarmati, *On the complexity of a family based on irreducible polynomials*, Preprint 2014.
5. ———, *On the complexity of a family related to the Legendre symbol*, Period. Math. Hungar. **58** (2009), no. 2, 209–215. MR 2531165 (2010f:11131)
6. ———, *Measures of pseudorandomness*, in Pascale Charpin, Alexander Pott, and Arne Winterhof (eds.), Finite Fields and Their Applications: Character sums and polynomials, Radon Ser. Comput. Appl. Math., vol. 11, Walter de Gruyter, Berlin, 2013, pp. 43–64.
7. Katalin Gyarmati, Christian Mauduit, and András Sárközy, *The cross correlation measure for families of binary sequences*, in Gerhard Larcher, Friedrich Pillichshammer, Arne Winterhof, and Chaoping Xing (eds.), Applications of Algebra and Number Theory, Cambridge University Press, to appear.
8. Henryk Iwaniec and Emmanuel Kowalski, *Analytic number theory*, American Mathematical Society Colloquium Publications, vol. 53, American Mathematical Society, Providence, RI, 2004. MR 2061214 (2005h:11005)
9. Christian Mauduit and András Sárközy, *On finite pseudorandom binary sequences. I. Measure of pseudorandomness, the Legendre symbol*, Acta Arith. **82** (1997), no. 4, 365–377. MR 1483689 (99g:11095)
10. Aimo Tietäväinen, *Vinogradov's method and some applications*, Number theory and its applications (Ankara, 1996), Lecture Notes in Pure and Appl. Math., vol. 204, Dekker, New York, 1999, pp. 261–282. MR 1661670 (99k:11186)
11. Alev Topuzoğlu and Arne Winterhof, *Pseudorandom sequences*, Topics in geometry, coding theory and cryptography, Algebr. Appl., vol. 6, Springer, Dordrecht, 2007, pp. 135–166. MR 2278037 (2007m:11106)
12. Arne Winterhof, *Some estimates for character sums and applications*, Des. Codes Cryptogr. **22** (2001), no. 2, 123–131. MR 1813781 (2002g:11128)

(Arne Winterhof, Oğuz Yayla) Johann Radon Institute for Computational and Applied Mathematics (RICAM), Austrian Academy of Sciences, Altenbergerstr. 69, 4040 Linz, Austria

E-mail address: arne.winterhof@ricam.oeaw.ac.at

E-mail address: oguz.yayla@ricam.oeaw.ac.at